

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 October 2001 (04.10.2001)


PCT

(10) International Publication Number
WO 01/72784 A2

- (51) International Patent Classification⁷: C07K 14/415, C12N 15/29, A61K 38/16, A61P 35/00
- (21) International Application Number: PCT/US01/09453
- (22) International Filing Date: 23 March 2001 (23.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/534,705 24 March 2000 (24.03.2000) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/534,705 (CIP)
Filed on 24 March 2000 (24.03.2000)
- (71) Applicant (for all designated States except US): FILGEN BIOSCIENCES, INC. [US/US]; 2333 Courage Drive, Suite C, Fairfield, CA 94533 (US).
- (72) Inventor; and
(75) Inventor/Applicant (for US only): GALVEZ, Alfredo, F. [PH/US]; 2800 Dover Avenue #G-5, Fairfield, CA 94533 (US).
- (74) Agents: PETERS, Howard, M. et al.; Peters, Verny, Jones & Biksa, LLP, Suite 6, 385 Sherman Avenue, Palo Alto, CA 94306 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: THERAPEUTIC PEPTIDES HAVING A MOTIF THAT BINDS SPECIFICALLY TO NON-ACETYLATED H3 AND H4 HISTONES FOR CANCER THERAPY



(57) Abstract: The present invention describes a composition of matter comprising a conserved structural motif that allows the targeting and binding of a chromatin binding protein to non-acetylated histone H3 and H4 and prevents their acetylation. This invention is responsible for the anti-carcinogenic property of a chromatin binding peptide isolated from soybean seed. This structural motif is found in a highly conserved manner in other chromatin-binding proteins from different species. Modifications to this structural motif such as fusions to other proteins with functional motifs and amino acid substitutions have potential therapeutic applications and can be developed as an *in vivo* gene silencing technology for biological and medical research. In particular, active fragments of the lunasin peptide and active analogs of the lunasin peptide are useful in this invention. Pharmaceutical compositions useful in retarding or stopping or reducing various types of cancers are described.

BEST AVAILABLE COPY

WO 01/72784 A2



Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

THERAPEUTIC PEPTIDES HAVING A MOTIF THAT BINDS
SPECIFICALLY TO NON-ACETYLATED H3 AND H4 HISTONES
FOR CANCER THERAPY

BACKGROUND OF THE INVENTION

5

Related Applications

This application is a continuation-in-part of U.S. Serial No. 09/534,705, filed March 24, 2000, which is incorporated herein by reference in its entirety.

10

Field of the Invention

This invention relates to lunasin, its fragments, analogs and the like which have a defined helical moiety which comprises a structurally conserved helical motif, a stretch of polyacidic amino acids (either aspartic or glutamic acid) and an Arg-Gly-Asp (RGD) for lunasin targeting and binding to non-acetylated N-terminal tails of H3 and H4 histones, making them unavailable for acetylation, and for cell membrane adherence and internalization. The substances are useful in a variety of disease therapy including reduction/repression of existing cancer or prevention of cancer initiation.

Description of Related Art

20

Lunasin, the small subunit of a soybean 2S albumin, colocalizes with endoreduplicated genomic has DNA in storage cells of developing seed.

25

The lunasin peptide with its unique poly-apartic acid carboxyl end and was proposed to have an important biological function when it was isolated and sequenced but not cloned from soybean seeds by a Japanese group 13 years ago (Odani et al., 1987 *J Biol Chem*, vol 262:10502). However, only upon the isolation and cloning of the *Gm2S-1* cDNA could a putative biological role for lunasin be inferred. The *Gm2S-1* cDNA encodes lunasin as a 43 amino acid small subunit component of a post-translationally processed 2S albumin (Galvez et al., 1997 *Plant Physiol*, vol. 114:1567). *Gm2S-1* expression occurs only in the cotyledon and coincides with the initiation of mitotic arrest and DNA endoreduplication in developing soybean seed (Galvez et al., 1997). DNA endoreduplication is a unique cell cycle of G1 and S phases without cell division that occurs only in terminally differentiated storage parenchyma cells (Goldberg et al, 1994 *Science*, vol. 266:605). *In situ* hybridization experiments using a

30

lunasin antisense RNA probe and immunolocalization using a polyclonal antibody raised against the carboxyl end of lunasin, showed lunasin expression in storage parenchyma cells undergoing DNA endoreduplication and cell expansion but not in actively dividing cells of the cotyledon (Fig. 1A, 1B, 1C, 1D and 1E.).

The temporal and spatial expression of lunasin in developing seeds suggest a biological role of lunasin as an effector molecule that inhibits cell division and allows DNA endoreduplication and cell expansion to occur in storage parenchyma cells during seed development. Its colocalization with endoreduplicated genomic DNA suggests a potential role as a repressor of gene expression in newly replicated genomic DNA. Despite the presence of multiple copies of the genome, the level of gene expression in storage parenchyma cells corresponds to a single copy of the genome. By binding to hypoacetylated chromatin associated with newly replicated DNA, lunasin is thought to silence expression of genes in the reduplicated genome by forming repressed chromatin structures. In addition, lunasin binding to hypoacetylated chromatin could inhibit mitotic condensation of the chromosomes and prevent microtubule nucleation, leading to the failure of cell division in expanding storage parenchyma cells. In support of this hypothesis, studies have shown that the phosphorylation of serine 10 in the amino terminal tail of histone H3 is required for the proper segregation and condensation of chromosomes during mitosis (Wei et al., 1999 *Cell*, vol. 97:99). Lunasin as described below has preferential binding affinity to the non-acetylated amino terminal tails of histone H3 and H4. By making the serine 10 unavailable for phosphorylation as a result of lunasin binding to the H3 amino terminal tail, lunasin can prevent condensation of chromosomes and consequently inhibit cell division. Constitutive expression of lunasin in mammalian cells

Disrupt centromere assembly and mitosis.

The temporal and spatial expression of lunasin coincide with the initiation of mitotic arrest and DNA endoreduplication in developing soybean cotyledon. This information, together with the observation that *lunasin* expression caused aberrant cell division in bacteria (Galvez and de Lumen, 1999 *Nature Biotechnology*, vol. 17:495), led to the hypothesis that lunasin should also disrupt eukaryotic cell division. To test this hypothesis, a chimeric gene encoding the lunasin peptide tagged with green fluorescent protein (GFP) was constructed. The

transient transfection of the GFP-lunasin construct arrested cell division, caused abnormal spindle fiber elongation, chromosomal fragmentation and cell lysis in murine embryo fibroblast, murine hepatoma, and human breast cancer cells (Galvez and de Lumen, 1999). Transfection of a control construct with a deleted poly-aspartyl end abolished lunasin's antimitotic effect.

5 The mechanism of action of other antimitotic agents such as vinblastine, colchicine, nocodazole and taxol involves the disruption of mitotic spindle dynamics during mitosis. Unlike these compounds, lunasin disrupts mitosis in mammalian cells by binding to chromatin and preventing the formation of the kinetochore complex in the centromere. This is likely brought about by the binding of the negatively charged lunasin to the highly basic histones
10 found within the nucleosomes of condensed chromosomes, particularly to regions that contain more positively charged, hypo-acetylated chromatin such as found in telomeres and centromeres. The displacement by lunasin of the kinetochore proteins normally bound to the centromere leads to the failure of spindle fiber attachment, and eventually to mitotic arrest and cell death. The observations of lunasin adhering to the fragmenting chromosomes after cell
15 lysis, the asymmetric distribution of metaphase chromosomes, the elongated spindle fibers, and the unattached kinetochores observed in *lunasin*-transfected cells are consistent with this proposed model for the mechanism of action of lunasin (Galvez and de Lumen, 1999).

Lunasin peptide adheres to mammalian cell membrane, gets internalized and binds to regions of hypoacetylated chromatin (i.e. telomeres)

20 Lunasin contains the cell adhesion motif RGD (arg-gly-aspartic acid). Synthetic and recombinant peptides containing the RGD motif derived from sequences of extracellular matrix proteins like fibronectin, have been shown to bind to specific membrane integrins in mammalian cells (E. Ruoslahti, M.D. Piersbacher. *Cell*, vol. 44, 517 (1986); S.K. Akiyama, K. Olden, K.M. Yamada. *Cancer Metastasis Rev.*, vol. 14, 173 (1995)). To determine whether lunasin has a
25 functional RGD motif, a cell adhesion assay using synthetic lunasin peptides and mice embryo fibroblast cells (C3H 10T1/2) was conducted (L.M. De Luca, *et al.*, *Methods of Enzymol.*, vol. 190:81-91 (1990)). The lunasin peptide adhered to C3H cells in a dose-dependent manner and that the deletion of the RGD tripeptide from lunasin (Lunasin-GRG) prevented cell

adhesion (Fig. 2). When applied exogenously to the growth media, lunasin was not only adhering to the cell membrane but became internalized as well, preferentially binding to the telomeres of chromosomes during metaphase (Fig. 3A, 3B, 3C, 3D, 3E, and 3F). However, unlike the constitutive expression of lunasin gene in transfected cells that disrupts kinetochore formation (Galvez and de Lumen, 1999), internalized lunasin did not affect kinetochore assembly. Immunostaining experiments showed the normal kinetochore location of the cell cycle checkpoint protein, MAD (Y. Li; R. Benezra, *Science*, vol. 274, 246 (1996); R.H. Chen, J.C. Waters, E.D. Salmon, A.W. Murray, *Science*, vol. 274, 242 (1996)), in the centromere of metaphase chromosomes. As a result, the exogenous application of lunasin did not affect cell division and proliferation of murine embryo fibroblast cells. Immunostaining using the lunasin polyclonal antibody also showed that internalized lunasin was initially found in the cytoplasm and then eventually bound to hypoacetylated regions of the chromosome, such as those in the telomeres, upon nuclear membrane breakdown at prometaphase (Fig. 3A, 3B, 3C, 3D, 3E, and 3F). However, at this stage of mitosis, kinetochore assembly and spindle fiber attachment to centromeres had already transpired. This explains the non-disruptive effect of exogenously applied lunasin on cell division as compared to the antimitotic effect observed when lunasin is constitutively expressed in *lunasin*-transfected mammalian cells (Galvez and de Lumen, 1999).

A U.S. patent of interest is U.S. 6,107,287 issued August 23, 2000.

All articles, references, standards, patents, patent applications and the like cited in this application are hereby incorporated herein by reference in their entirety.

With regard to the above background description, there exists a significant need to provide a method and pharmaceutical composition to inhibit or retard various cancers from initializing and or reducing existing cancers for shrinking particularly in a human being. The present invention provides such a method and pharmaceutical composition.

SUMMARY OF THE INVENTION

The present invention relates to a method of cancer treatment or prevention, which method involves:

A. Administering to a mammalian subject having tumor cells in need of therapy or

a mammalian subject at risk to carcinogen-mediated cancer formation an effective amount of an isolated and purified therapeutic agent selected from the group consisting of lunasin peptide, an active fragment of lunasin peptide, an active lunasin peptide analog and combinations thereof which lunasin moiety has a helical portion which comprises the structural motif (ED)NNXXXEK(IV), where E is glutamic acid, D is aspartic acid, K is lysine, I is isoleucine, V is valine, X is conserved hydrophobic amino acids and N is any amino acid, a sequence of at least 5 up to 15 poly-acidic amino acids (glutamic or aspartic acids), and an Arg-Gly-Asp (RGD) motif which is useful for targeting and binding to non-acetylated N-terminal tails of H4 and H3 histones and for functional adhesion of lunasin moiety to the outer cell membrane;

10 B. Causing the lunasin peptide, the active fragment of lunasin peptide, the active lunasin peptide analog or combinations thereof to contact and to adhere to the functional cell membrane;

15 C. Causing the lunasin peptide, the active fragment of lunasin peptide, the active lunasin peptide analog or combinations thereof to contact and to become internalized within the functioning cell;

20 D. Causing the lunasin peptide, the active fragment of lunasin peptide, the active lunasin peptide analog or combinations thereof to preferentially bind to the deacetylated N-terminal portions of histone H3 and H4, causing these histones to be unavailable for further acylation in regions of the chromosomes of the cell and which are enriched with hypoacetylated repressed chromatin;

E. Inducing apoptosis of the cell by repression of carcinogen and oncogene-mediated gene expression within the cell; and

F. Resulting in significantly reduced or termination of cancer activity of existing tumor cells or the prevention of significant tumor cell initiation.

25 The method wherein the mammal is a human being.

The method wherein the method is one of treating an already existing cancer.

30 The method wherein the method is one of preventing or repressing the induction of cancer.

The method wherein the therapeutic agent comprises lunasin peptide.

The method wherein the therapeutic agent comprises an active fragment of lunasin peptide.

5 The method wherein the therapeutic agent comprises an active analog of lunasin peptide.

The method wherein the therapeutic agent is administered orally, topically, intranasally, intramuscularly, subcutaneously, intraperitoneally, buccally or combinations of these methods.

10 The method wherein the therapeutic agent is administered topically in a pharmaceutically acceptable excipient.

In another aspect the present invention concerns a method and a pharmaceutical composition wherein the pharmaceutical composition is administered topically to retard or stop cancers of the skin.

15 In another aspect the present invention concerns a method and a pharmaceutical composition wherein the pharmaceutical composition is administered intranasally or as part of inhalation therapy to retard or stop cancers of the lung.

In another aspect the present invention concerns a method and a pharmaceutical composition wherein the pharmaceutical composition is administered intravenously to retard or stop cancers of the breast, prostate, liver, kidney or any other internal organs or tissues.

20 In another aspect the present invention concerns a method and a pharmaceutical composition wherein the pharmaceutical composition is administered is a vaginal suppository to retard or stop cancers of the cervix, uterus or ovary.

25 In another aspect the present invention concerns a method and a pharmaceutical composition wherein the pharmaceutical composition is administered as an anally applied suppository to retard or stop cancers of the lower gastro-intestinal tract.

In another aspect the present invention concerns a method and a pharmaceutical composition wherein the pharmaceutical composition is administered orally to retard or stop cancers of the colon, upper gastrointestinal tract, breast, prostate, liver, kidney or any other internal organs or tissues.

30 In another aspect the present invention concerns a method and a pharmaceutical composition wherein the pharmaceutical composition is administered intramuscularly or

subcutaneously as a general protection against cancer development in internal organs.

In another aspect, the present invention concerns a method of targeting and binding non-acetylated H3, H4 histones and other histone -variants such as the centromere-specific H3 variant, CENP-A, which method comprises:

- 5 A. Prevention of acetylation of amino acid residues found in N-terminal tail of H3, H4 and variant histones,
- B. Prevention of phosphorylation of amino acid residues found in N-terminal tails of H3, H4 and variant histones.
- C. Prevention of methylation of amino acid residues found in N-terminal tails of H3, H4 and variant histones.
- 10 D. Prevention of other post-translational modifications of amino acid residues found in N-terminal tails of H3, H4 and variant histones, with the result.

In another aspect, the present invention concerns a composition of matter, that is required to allow targeting and binding of proteins to non-acetylated H3, H4 histones and other histone -variants such as the centromere-specified H3 variant, CENP-A, which composition
15 comprises:

- A. Presence of a helical motif that is structurally conserved, comprising a consensus sequence of 9 amino acid residues, composed of: (ED)NNXXXEK(IV), where E is glutamic acid, D is aspartic acid, I is isoleucine, V is valine, K is lysine residues, N is any amino acid, and X is conserved hydrophobic residues, and the
- 20 B. Presence of a block of 5-10 residues of acidic amino acids (either E is glutamic acid or D is aspartic acid), upstream or downstream of the helical motif.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A, 1B, 1C, 1D and 1E are schematic representations of lunasin found in storage
25 parenchyma cells and co-localizes with endoreduplicated DNA.

Figure 2 is a graphic representation of relative cell adhesion versus amount of peptide added for lunasin and lunasin (-GRC) as it attaches to mammalian cell membrane through its RGD motif.

Figures 3A, 3B, 3C, 3D, 3E and 3F are schematic representations of lunasin adhering to

the cell membrane and then becoming internalized.

Figures 4A and 4B are schematic representations of lunasin as a major constituent of the Bowman Birk protease inhibitor (BBIC) preparation.

Figure 5 is a graphic representation of how lunasin inhibits carcinogen-induced transformation.

5 Figure 6 is a graphic representation of lunasin in prevention of carcinogen-induced tumorous foci formation in normal cells.

Figures 7A, 7B, 7C, 7D, 7E and 7F are photographic representations of C3H cells transfected with E1A- Δ CR1, in the absence of lunasin and the presence of lunasin which induces apoptosis.

10 Figure 8 is a schematic representation and model for the prevention of cancer in the presence of lunasin.

Figure 9 is a graphic representation showing lunasin preferentially binding to deacetylated histone H4.

15 Figure 10 is a graphic representation showing the dose response of lunasin, trLunasin-del and NLS-trLunasin to increasing amounts of deacetylated H4 peptide.

Figure 11 is a table which compares motifs showing that lunasin contains a helical motif having high structural homology to other chromatin binding proteins.

Figure 12 is a graphic representation of the effect of modified lunasin peptides on transformation assay.

20 Figure 13 is a schematic representation depicting how lunasin binds to deacetylated histones and inhibits histone acylation.

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

25 Definitions

As used herein:

The standard literature definitions found in articles and reference books are to be used to determine the definitions of the terms as found herein.

30 "Amino acid" refers to any of the naturally occurring amino acids having standard designations, G, V, K, I, W, etc. It also refers to those known synthetic amino acids.

Conserved hydrophobic amino acid refers to but are not limited to, for example, histidine, isoleucine, valine, methionine, alanine, or tyrosine.

“Lunasin” refers to compounds comprising the natural and recombinantly produced soybean lunasin polypeptide (coincidentally purified and sequenced by Odani et al., 1987 (Ser-
5 Lys-Trp-Gln-His-Gln-Gln-Asp-Ser-Cys-Arg-Lys-Gln-Leu-Gln-Gly-Val-Asn-Leu-Thr-Pro-Cys-Glu-Lys-His-Ile-Met-Glu-Lys-Ile-Gln-Gly-Arg-Gly-Asp-Asp-Asp-Asp-Asp-Asp-Asp-Asp (SEQ. ID. 1).

“Lunasin” refers to the biologically active lunasin peptide having 1-43 amino acids.

“Lunasin or an active variant thereof” refers to the biologically active lunasin peptide
10 having 43 amino acids, or to portions of the 1-43 amino acid chain which are also biologically active (shown herein as 22-43 amino acids meaning amino acid 22 to amino acid 43 of lunasin). See sequence data below:

protein having amino acids 1 to 42 (SEQ. ID.2),
protein having amino acids 1 to 41 (SEQ. ID.3),
15 protein having amino acids 1 to 40 (SEQ. ID.4),
protein having amino acids 1 to 39 (SEQ. ID.5),
protein having amino acids 1 to 38 (SEQ. ID.6).
protein having amino acids 22 to 43 (SEQ. ID.7),
protein having amino acids 22 to 42 (SEQ. ID.8),
20 protein having amino acids 22 to 41 (SEQ. ID.9),
protein having amino acids 22 to 40 (SEQ. ID.10),
protein having amino acids 22 to 39 (SEQ. ID.11), and
protein having amino acids 22 to 38 (SEQ. ID.12)

Combinations of these active protein are also included.

25 Polyacidic amino acids refer, for example, to glutamic acid or aspartic acid.

1. The lunasin peptide has anti-carcinogenic property

Lunasin has been shown to be a major constituent of the Bowman Birk protease inhibitor (BBIC) preparation (Fig. 4A and 4B). BBIC has been shown to be chemopreventive in several *in vitro* and animal model studies (Examples: Yavelow et al., 1985 *PNAS*, vol.

82:5395; Weed et al., 1985 *Carcinogenesis*, vol 6:1239; Messadi et al., 1986 *JNCI*, vol. 76:447; Baturay, et al., 1986 *Cell Biol and Toxic*, vol. 2:21; St. Clair et. al., 1990 *Cancer Res*, vol 50:580; Reviews: Kennedy et al., 1993 *Preventive Med*, vol 22:796; Kennedy et al., 1995 *J Nutr*, vol. 125:733S). The evidence for the anti-carcinogenic effect of BBIC was compelling enough that NCI is now conducting human clinical trials (currently in Phase II) to prove its effectivity (Kennedy et al., 1993 *Preventive Med*, vol. 22:796). However, despite the accumulated *in vitro* and *in vivo* data pointing to the anticarcinogenic property of BBIC, the underlying mechanism of action has not been elucidated. More importantly, several scientific evidence have shown that BBIC or protease inhibitors (PI), in general, are unlikely to be the active anticarcinogenic component found in soybean. For one, cooked soy products, which are devoid of any protease inhibitor activity, are equally as effective at reducing cancer development as raw soy products (Clawson, 1996 *Cancer Invest.*, vol. 14(6):608). The effect of protease inhibitors appears to be indirect because dietary PI are, in general, poorly absorbed from the gastro-intestinal (GI) tract, and never reach target organs in any measurable quantity (Clawson, 1996).

Lunasin is responsible for the cancer preventive activity attributed to BBIC, specially since the lunasin peptide is a significant contaminant in the BBIC preparation. Cell transformation assays conducted at UC Berkeley showed that lunasin was on average twice more effective than equimolar amounts (125 nM) of BBIC in reducing foci formation in C3H 10 T1/2 cells treated with potent chemical carcinogens, 7, 12-dimethylbenz[a]anthracene (DMBA) and 3-methylcholanthrene (MCA) (Fig.5). More importantly, BBIC with immunodepleted lunasin, prepared by applying commercially available BBI (Sigma T9777) through cationic exchange and immuno-affinity columns and then collecting flow through fractions, showed significant loss of its anti-transformation property (Fig.6). The duplicated sets of experiments showed that BBIC with immunodepleted lunasin did not inhibit foci formation upon carcinogen treatment, similar to the effect of the untreated positive control. These results indicate that lunasin is the major cancer preventive ingredient in the BBIC preparation.

What then is the role of BBI in the cancer preventive property attributed to the BBIC soybean preparation? As pointed out by Clawson (1996), the effect of BBI appears to be

indirect. Digestion experiments have shown that lunasin by itself gets broken down by pancreatic digestive enzymes but resists digestion when a chymotrypsin inhibitor like BBI is mixed with lunasin at equimolar ratios (Pascual and de Lumen, personal communication). It is most likely that BBI's role is to prevent the digestion of lunasin in the gut to allow intact lunasin to be absorbed through the gastro-intestinal tract. Once in the circulatory system, lunasin can be distributed to the various tissues and can get inside somatic cells by attaching to specific integrin receptors found in cell membranes through its RGD cell adhesion motif. Inside the cell, lunasin then preferentially binds to regions of the chromosomes enriched with hypoacetylated chromatin upon nuclear membrane breakdown at prometaphase.

2. Anti-carcinogenic property of lunasin: a molecular model based on lunasin binding to deacetylated histones and inhibition of histone acetylation.

The affinity of the lunasin peptide to regions of hypoacetylated chromatin suggests that lunasin may be involved in chromatin modification. Regulation of the post-translational modification of chromatin has been implicated in cell-cycle control and in how tumor suppressors act as critical downstream effectors during carcinogenesis (R.A. DePinho, *Nature*, vol. 391, 533 (1998)). Lunasin also contains a functional cell adhesion motif, Arg-Gly-Asp (RGD), which allows exogenously applied lunasin to bind and become internalized in mammalian cells. The presence of the RGD motif and its chromatin-binding characteristic point to a potential anti-carcinogenic role for lunasin.

Histone acetylation is associated with transcriptional activity in eukaryotic cells, having been observed mainly in transcriptionally active chromatin (K. Struhl, *Genes Dev.*, vol. 12, 599 (1998); M. Grunstein, *Nature*, vol. 389, 349 (1997)). The inhibition of histone acetylation by lunasin provides a mechanistic model to explain the anti-carcinogenesis property of this soybean peptide. The Rb tumor suppressor, a critical downstream effector during carcinogenesis (R.A. Weinberg, *Cell*, vol. 81, 323 (1995); M.C. Paggi, *et al.*, *J Cell. Biochem.*, vol. 62, 418 (1996)), was hypothesized to repress a subset of E2F-regulated genes by binding to the E2F family of DNA-binding transcription factors and by recruiting a histone deacetylase (HDAC1) to maintain a hypoacetylated state of condensed chromatin around the transcription start site (A. Brehm *et al.*, *Nature*, vol. 391, 597 (1998); L. Managhi-Jaulin *et al.*, *Nature*, vol. 391, 601 (1998); R.X. Luo, A.A. Postigo, D.C. Dean, *Cell*, vol. 92, 463

(1998)). This dual repression mechanism is abrogated upon Rb inactivation during carcinogenesis, resulting in the release of Rb binding to the E2F promoter, acetylation of the repressed chromatin structure and the induction of expression of the E2F-regulated genes involved in cell proliferation (A. Brehm *et al.*, *Nature*, vol. 391, 597 (1998); L. Managhi-Jaulin *et al.* *Nature*, vol. 391, 601 (1998); R.X. Luo, A.A. Postigo, D.C. Dean, *Cell*, vol. 92, 463 (1998)).

By binding to deacetylated histones found in repressed chromatin, it was hypothesized that lunasin can prevent cell proliferation and transformation even in the absence of a functional Rb by inhibiting histone acetylation and activation of E2F-regulated genes. To test this molecular model of lunasin action, C3H cells were first treated with lunasin and then transfected with E1A viral oncogene that specifically induces cell proliferation by binding and inactivating Rb (J.R. Nevins, *Science*, vol. 258, 424 (1992)). As a negative control, E1A with deleted conserved region 1 (E1A Δ CR1) that abolishes the RB binding domain was likewise used in the transfection experiments (D. Trouche, T. Kouzidares, *Proc. Natl. Acad. Sci., USA*, vol. 93, 1439 (1996)). C3H cells transfected with E1A- Δ CR1, as expected, showed normally dividing cells at 20 h after transfection, both in the presence and absence of lunasin (Fig. 7A, 7B, 7C, 7D, 7E and 7F). Transfection with the E1Awt in the absence of lunasin also showed normal cell proliferation (Fig. 7A, 7B, 7C, 7D, 7E and 7F). However, C3H cells initially treated with lunasin for 24h and then transfected with E1Awt resulted in the preponderance of non-adherent cells in solution at 20 h after transfection. Phase contrast image of the non-adherent cells showed characteristic morphology of apoptotic cells which was confirmed by the positive fluorescent staining for Annexin V-FITC (Fig. 7A, 7B, 7C, 7D, 7E and 7F.).

The induction of apoptosis by lunasin in E1A-transfected C3H cells provides evidence to a mechanistic model explaining lunasin's suppression of carcinogen-mediated transformation (Fig.8). The Rb tumor suppressor inhibits the expression of E2F-regulated genes in part by tethering a histone deacetylase (HDAC1) to maintain a condensed hypoacetylated chromatin around the transcription start site (A. Brehm *et al.*, *Nature*, vol. 391, 597 (1998); L. Managhi-Jaulin *et al.* *Nature*, vol. 391, 601 (1998); R.X. Luo, A.A. Postigo, D.C. Dean, *Cell*, vol. 92, 463 (1998)). The inactivation of Rb by carcinogen treatment and oncogene expression results in the loosening up of the repressed chromatin structure by

localized histone acetylation (R.H. Giles, D.J. Peters, M.H. Breuning, *Trends Genet.*, vol. 14, 178 (1998)). This consequently results in the activation of genes involved in cell proliferation, which eventually leads to carcinogenesis. When lunasin is present in normal cells before Rb is inactivated, the deacetylated N-terminal tails of histone H3 and H4 found in repressed chromatin presumably bind to the acidic carboxyl end of lunasin. This makes these deacetylated histones unavailable as substrates for histone acetylation, thus maintaining the repressed chromatin structure around the E2F promoter even when carcinogens and the viral oncogene, E1A, inactivate Rb. The inhibition of expression of E2F-regulated genes triggers apoptosis instead of cell proliferation, which normally occurs when these genes are activated during carcinogenesis. The induction of apoptosis in cells with inactivated Rb by the presence of lunasin can explain the reduced number of transformed foci in normal murine fibroblast cells that have been treated with potent chemical carcinogens.

UTILITY AND ADMINISTRATION - Administration of the compounds of this invention can be via any of the accepted modes of administration for therapeutic agents. These methods include oral, parenteral, transdermal, subcutaneous and other modes.

Depending on the intended mode, the composition may be in many forms, for example, solid, semi-solid, or liquid dosage forms, including tablets, time release agents, pills, capsules, suspensions, solutions and the like. The compositions will include a conventional pharmaceutical excipient and an active compound as described herein or the pharmaceutically acceptable salts thereof and may, in addition, include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, etc.

The amount of the active compound administered will, of course, be dependent on the molecular weight of selected compound, the subject being treated, the subject's weight, the severity of the affliction, the manner of the administration and the judgment of the prescribing physician. However, an effective dose is in the range of about 0.1-500 mg/kg/day, preferably about 1-200 mg/kg/day. For an average 70 kg human, those dosages would amount to between about 0.01 to 35 g/day.

For solid compositions, conventional nontoxic solids include for example, pharmaceutical grades of manitol, lactose, starch, magnesium stearate, cellulose and the like may be used. Liquid pharmaceutically administratable compositions can be prepared by dissolving,

dispersing, etc., a compound and optional pharmaceutical adjuvants in an excipient, such as, for example, water, glycerol, ethanol, vegetable oil and the like to form a suspension.

Actual methods of preparing such dosage forms are known, or will be apparent to those skilled in the art; see, for example, *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Easton, Penn., 15th Edition, 1975.

5 For instance for topical or intranasal or intravenous administration, the minimum dose is about 250 microg lunasin per mL of solution or per gram of solid dose up to a maximum dose of about 2.5 millig lunasin per mL of solution or per gram of solid dose.

The following preparations and examples serve to illustrate the invention. They should not be construed as narrowing it, nor as limiting its scope.

10 Experimental - General

The starting materials described herein are available from commercial supply houses, from recognized contracting organizations or can be prepared from published literature sources. Unless otherwise noted the material solvents, reagents, etc. are used as received without modification.

15 EMBODIMENTS OF THE INVENTION:

The experimental evidence described above point to the utility of the lunasin peptide in disrupting specific cellular processes like carcinogenesis. The proposed lunasin mechanism of action involves its preferential binding to the deacetylated N-terminal tails of histone H3 and H4, making them unavailable as substrates for acetylation. Since the acetylation of histone H3 and H4 is associated with gene activation, lunasin acts as a repressor of gene expression when it binds to deacetylated histones found in promoter regions of negatively regulated genes (such as the family of E2F-regulated genes that are negatively regulated by the Rb tumor suppressor). The ability of lunasin to repress gene expression by preferential binding to deacetylated histones and preventing their acetylation has practical wide-ranging biological and therapeutic applications.

25 The invention describes the identification of the functional motif in the lunasin peptide responsible for its chromatin-binding property and its ability to inhibit acetylation of H3 and H4 histones. This invention is important for designing future drugs involving targeted repression of genes and for practical application in biological research by providing a method

to target modified lunasin peptides to specific genes or genome locations and for the study of phenotypic effects of gene inactivation and silencing.

EXAMPLE 1

BINDING OF LUNASIN AND FRAGMENTS THEREOF

- 5 (a) The lunasin peptide preferentially binds to deacetylated histones and is mediated by a helical region in the carboxyl end.

The antimitotic effect of the lunasin gene in transfected mammalian cells has been attributed to the competitive binding of lunasin to centromeres as visualized by GFP fluorescence and immunostaining (Galvez and de Lumen, 1999). On the other hand,
10 immunostaining of exogenously applied lunasin revealed the preferential binding of lunasin mainly to the telomeres of metaphase chromosomes (Fig. 3). Telomeres, like the centromeres are genomic regions that are also rich in hypoacetylated chromatin, comprising mainly of deacetylated histones (Braunstein *et al.*, *Genes Dev*, vol. 7, 592, 1993). The increased affinity of lunasin to these regions may be due to the greater electrostatic attraction of the negatively
15 charged carboxyl end of lunasin to the positively charged N-terminal tails of deacetylated histones.

To test whether lunasin binds preferentially to deacetylated histones, an *in vitro* immunobinding assay was conducted using acetylated and deacetylated forms of the H4 N-terminal tail (assay protocol was described in Galvez and de Lumen, 1999). The full lunasin peptide
20 (Lunasin) and lunasin with deleted RGD motif (Lunasin-GRG) were found to bind with high affinity to deacetylated H4 N-terminus but not to the tetra-acetylated H4 (Fig. 9). This suggests that lunasin binds with high specificity to deacetylated H4 and that the RGD-motif is not important to its binding affinity. However, there was a significant reduction in deacetylated H4 binding for truncated lunasin (trLunasin) that contains only the reactive
25 carboxyl end of the peptide. This indicates that the N-terminus of lunasin is also important for binding to deacetylated histones most likely by stabilizing the lunasin structure to allow electrostatic interactions between the carboxyl end of lunasin and deacetylated H4 to occur at higher efficiency.

A comparison of the binding affinity of lunasin, tr-Lunasin and NLS-trLunasin to

increasing dose of deacetylated H4 peptide showed an increase in lunasin binding when the amount of deacetylated H4 peptide added in the immuno-binding reaction is increased (Fig. 10). Lunasin binding to deacetylated H4 was 3X more than trLunasin-del, which in return was found to bind at significantly higher affinity than the NLS-trLunasin (Fig. 10).

The binding affinity of trLunasin to deacetylated H4 was not significantly different from that of the 10 amino acid trLunasin-del peptide fragment. The trLunasin-del fragment spans a helical domain (B. Rost, C. Sander, *Proteins*, vol. 19, 55 (1994); B. Rost, C. Sander, *J Mol. Biol.*, vol. 232, 584 (1993)) upstream of the poly-aspartyl carboxyl end of the lunasin peptide. The substitution of this helix by a nuclear localization sequence (NLS) in the truncated lunasin peptide (NLS-trLunasin) resulted in the loss of binding to deacetylated H4 (Fig. 9). This indicates that this helical region of lunasin may play a role in the binding of lunasin to deacetylated histones. A homology search of this helical region revealed structural similarity to a short, conserved region of the chromo-domain structure (R. Aasland, A.F. Stewart, *Nucleic Acids Res*, vol. 23, 3168 (1995)) found in chromatin-binding proteins such as *Drosophila* and human heterochromatin (DmHP1A and HuHP1B, respectively) (Fig. 11). A naturally occurring mutation in *Drosophila* DmHP1A that converts isoleucine to phenylalanine (I to F mutation) (Fig. 11) led to the disruption of the helical motif and the consequent loss of chromatin targeting (S. Messmer, A. Franke, R. Paro, *Genes Dev.*, vol. 6, 1241 (1992)). The presence of this helical motif in lunasin could explain the specific targeting of the peptide to deacetylated chromatin. Its absence from the NLS-trLunasin peptide reduced the binding to deacetylated H4 significantly, despite the presence of the poly-aspartyl end (Fig. 9). However, the presence of both helix and poly-aspartyl end was necessary for binding to deacetylated H4 (Fig. 9) and for the anti-transformation property of the truncated lunasin (trLunasin) peptide (Fig. 11). The poly-aspartyl end attached to this helical motif at the carboxyl end appears to be important for the anti-carcinogenic property of lunasin. Although the helical motif is necessary for targeting the lunasin peptide to deacetylated histones, it is the acidic poly-aspartyl end that interacts with the positively charged non-acetylated lysine residues in the histone N-terminal tails preventing them from being acetylated. It should also be pointed out that trLunasin has a lower binding affinity to deacetylated H4 than the full-length lunasin peptide (Fig. 9). This observation correlates with

the reduced efficacy of trLunasin in preventing foci transformation (Fig. 12). This result provides evidence linking the binding affinity of lunasin to deacetylated histones and its anti-transformation property in vivo, preferably in a human being.

(b) Similarly when the reaction involving lunasin (SEQ.ID.1 of 43 amino acids) of step (a) is repeated except that the lunasin is replaced by a stoichiometrically equivalent and active

5 fragment selected from:

protein having amino acids 1 to 42 (SEQ. ID.2),

protein having amino acids 1 to 41 (SEQ. ID.3),

protein having amino acids 1 to 40 (SEQ. ID.4),

protein having amino acids 1 to 39 (SEQ. ID.5),

10 protein having amino acids 1 to 38 (SEQ. ID.6).

protein having amino acids 22 to 43 (SEQ. ID.7),

protein having amino acids 22 to 42 (SEQ. ID.8),

protein having amino acids 22 to 41 (SEQ. ID.9),

protein having amino acids 22 to 40 (SEQ. ID.10),

15 protein having amino acids 22 to 39 (SEQ. ID.11), and

protein having amino acids 22 to 38 (SEQ. ID.12);

a corresponding useful therapeutic result is obtained in cancer inhibition and in reduction of cancer activity in vivo.

20

EXAMPLE 2

INHIBITION OF IN VIVO ACETYLATION

(a) Lunasin binding to deacetylated histones inhibits *in vivo* acetylation of histone H3 and H4

25 The *in vitro* binding of lunasin to deacetylated histone H4 confirms the observed affinity of lunasin to regions of hypoacetylated chromatin such as the centromeres and telomeres in immunostaining experiments (Galvez and de Lumen, 1999 and Fig. 6). Deacetylated histones are substrates for histone acetylation and for chromatin remodelling which has been associated with eukaryotic transcriptional regulatory mechanisms (K. Struhl, *Genes Dev.*, vol. 12, 599, 1998; M. Grunstein, *Nature*, vol. 389, 349, 1997). To determine whether the preferential

binding of lunasin to deacetylated histones has any biochemical effect on histone acetylation *in vivo*, C3H cells and the human breast cancer cell line, MCF-7, were treated with the histone deacetylase inhibitor, Na-butyrate (E.P. Candido, R. Reeves, J.R. Davie, *Cell*, vol. 14, 105, 1978), in the presence or absence of lunasin. Immunoblots of acid-extracted proteins show the significant reduction of acetylated H4 and H3 in Na-butyrate treated C3H and MCF-7 cells when pretreated with 1 μ M of lunasin peptide (Fig. 13). The absence of lunasin when cells were treated with Na-butyrate increased histone H4 acetylation by 200 fold in both C3H and MCF-7 cells. H3 acetylation induced by Na-butyrate treatment increased 100 fold in C3H cells and around 400 fold in MCF-7 cells. Upon addition of lunasin, there was no observed increase in H4 and H3 acetylation of C3H cells treated with Na-butyrate. In MCF-7 cells, H4 acetylation was reduced 10 fold and H3 acetylation 4 fold when lunasin was added prior to Na-butyrate treatment. These results demonstrate that the exogenous application of the lunasin peptide inhibit histone acetylation of mammalian cells *in vivo*, preferably in a human being.

(b) Similarly when the reaction involving lunasin (SEQ.ID.1 of 43 amino acids) of step (a) is repeated except that the lunasin is replaced by a stoichiometrically equivalent and active fragment selected from:

- protein having amino acids 1 to 42 (SEQ. ID.2),
- protein having amino acids 1 to 41 (SEQ. ID.3),
- protein having amino acids 1 to 40 (SEQ. ID.4),
- protein having amino acids 1 to 39 (SEQ. ID.5),
- protein having amino acids 1 to 38 (SEQ. ID.6).
- protein having amino acids 22 to 43 (SEQ. ID.7),
- protein having amino acids 22 to 42 (SEQ. ID.8),
- protein having amino acids 22 to 41 (SEQ. ID.9),
- protein having amino acids 22 to 40 (SEQ. ID.10),
- protein having amino acids 22 to 39 (SEQ. ID.11), and
- protein having amino acids 22 to 38 (SEQ. ID.12),

a corresponding useful therapeutic result is obtained in cancer inhibition and in reduction of existing cancer activity *in vivo*, preferably in a human being.

While only a few general embodiments of the invention have been shown and described herein, it will become apparent to those skilled in the art that various modifications and changes can be made in the application of lunasin and lunasin analogs and active lunasin fragments thereof to treat existing tumors or prevent initiation of tumor formation without departing from the spirit and scope of the present invention. All such modifications and

5 changes coming within the scope of the appended claims are intended to be carried out thereby.

I CLAIM:

1. A method of cancer treatment or prevention, which method comprises:
 - A. Administering to a mammalian subject having tumor cells in need of therapy or a mammalian subject at risk to carcinogen or oncogene-mediated cancer formation an effective amount of an isolated and purified therapeutic agent selected from the group consisting of lunasin peptide, an active fragment of lunasin peptide, an active lunasin peptide analog and combinations thereof which lunasin moiety has a helical portion which the structural motif (ED)NNXXXEK(IV), where E is glutamic acid, D is aspartic acid, K is lysine, I is isoleucine, V is valine, X is selected from conserved hydrophobic amino acids and N is any amino acid, a sequence of at least 5 to about 15 poly-acidic amino acids selected from glutamic acid or aspartic acid, and an Arg-Gly-Asp (RGD) motif which is useful for targeting and binding to non-acetylated N-terminal tails of H4 and H3 histones and for functional adhesion of lunasin moiety to the outer cell membrane;
 - B. Causing the lunasin peptide, the active fragment of lunasin peptide, the active lunasin peptide analog or combinations thereof to contact and to adhere to the functional cell membrane;
 - C. Causing the lunasin peptide, the active fragment of lunasin peptide, the active lunasin peptide analog or combinations thereof to become internalized within the functioning cell;
 - D. Causing the lunasin peptide, the active fragment of lunasin peptide, the active lunasin peptide analog or combinations thereof to preferentially bind to the deacylated N-terminal portions of histone H3 and H4, causing these histones to be unavailable for further acylation in regions of the chromosomes of the cell and which are enriched with hypoacylated repressed chromatin;
 - E. Inducing apoptosis of the cell by repression of carcinogen-mediated gene transformation within the cell, which results in significantly reduced or termination of cancer activity of existing tumor cells or the prevention of significant tumor cell initiation.

2. The method of claim 1 wherein the mammal is a human being.
3. The method of Claim 1 wherein the method is one of treating an already existing cancer.
4. The method of claim 1 wherein the method is one of preventing or repressing the induction of cancer.
- 5 5. The method of Claim 1 wherein the therapeutic agent comprises lunasin peptide.
6. The method of Claim 1 wherein the therapeutic agent comprises an active fragment of lunasin peptide.
7. The method of Claim 6 wherein the active fragment of lunasin is selected from the group consisting of:
 - 10 protein having amino acids 1 to 42 (SEQ. ID.2),
 - protein having amino acids 1 to 41 (SEQ. ID.3),
 - protein having amino acids 1 to 40 (SEQ. ID.4),
 - protein having amino acids 1 to 39 (SEQ. ID.5),
 - protein having amino acids 1 to 38 (SEQ. ID.6).
 - 15 protein having amino acids 22 to 43 (SEQ. ID.7),
 - protein having amino acids 22 to 42 (SEQ. ID.8),
 - protein having amino acids 22 to 41 (SEQ. ID.9),
 - protein having amino acids 22 to 40 (SEQ. ID.10),
 - protein having amino acids 22 to 39 (SEQ. ID.11),
 - 20 protein having amino acids 22 to 38 (SEQ. ID.12), and combinations thereof.
8. The method of Claim 1 wherein the therapeutic agent comprises an active analog of lunasin peptide.
9. The method of Claim 1 wherein the therapeutic dose is about 250 microgram per milliliter or per gram of solid dose to about 2.5 milligram per milliliter or per gram of solid
25 dose.

10. The method of Claim 1 wherein the therapeutic agent is administered orally, topically, intranasally, intramuscularly, subcutaneously, intraperitoneally, buccally intravenously or combinations of these methods.

11. The method of Claim 1 wherein the therapeutic agent is administered topically in a pharmaceutically acceptable excipient.

5 12. The method of Claim 1 wherein the therapeutic agent is administered orally.

13. A pharmaceutical composition which comprises a lunisin peptide, an active fragment of lunasin peptide, an active lunasin peptide analog or combinations thereof and a pharmaceutically acceptable excipient.

10 14. The pharmaceutical composition of Claim 13 which comprises a lunisin peptide and a pharmaceutically acceptable excipient.

15 15. The pharmaceutical composition of Claim 13 which comprises an active fragment of lunasin peptide, and a pharmaceutically acceptable excipient.

16. The pharmaceutical composition of Claim 13 wherein the active fragment of lunasin peptide is selected from the group consisting of:

- 15 protein having amino acids 1 to 42 (SEQ. ID.2),
protein having amino acids 1 to 41 (SEQ. ID.3),
protein having amino acids 1 to 40 (SEQ. ID.4),
protein having amino acids 1 to 39 (SEQ. ID.5),
protein having amino acids 1 to 38 (SEQ. ID.6).
20 protein having amino acids 22 to 43 (SEQ. ID.7),
protein having amino acids 22 to 42 (SEQ. ID.8),
protein having amino acids 22 to 41 (SEQ. ID.9),
protein having amino acids 22 to 40 (SEQ. ID.10),
protein having amino acids 22 to 39 (SEQ. ID.11),
25 protein having amino acids 22 to 38 (SEQ. ID.12), and combinations thereof.

17. The pharmaceutical composition of Claim 13 which comprises an active lunasin peptide analog and a pharmaceutically acceptable excipient.

18. The pharmaceutical composition of Claim 13 wherein the therapeutic dose is about 250 microg per milliliter or per gram of solid dose to about 2.5 millig per milliliter or per gram of solid dose.

19. The pharmaceutical composition of Claim 13 wherein said pharmaceutical composition is administered orally, topically, intranasally, intramuscularly, subcutaneously, 5 intrapertineally, buccally or combinations of these methods.

20. The pharmaceutical composition of Claim 13 wherein said pharmaceutical composition is administered topically to retard or stop cancers of the skin.

21. The pharmaceutical composition of Claim 13 wherein said pharmaceutical composition is administered intranasally or as part of inhalation therapy to retard or stop 10 cancers of the lung.

22. The pharmaceutical composition of Claim 13 wherein said pharmaceutical composition is administered intravenously to retard or stop cancers of the breast, prostate, liver, kidney or any other internal organs or tissues.

23. The pharmaceutical composition of Claim 13 wherein said pharmaceutical 15 composition is administered is a vaginal suppository to retard or stop cancers of the cervix, uterus or ovary.

24. The pharmaceutical composition of Claim 13 wherein said pharmaceutical composition is administered as an anally applied suppository to retard or stop cancers of the lower gastro-intestinal tract.

20 25. The pharmaceutical composition of Claim 13 wherein said pharmaceutical composition is administered orally to retard or stop cancers of the colon, upper gastrointestinal tract, breast, prostate, liver, kidney or any other internal organs or tissues.

26. The pharmaceutical composition of Claim 13 wherein said pharmaceutical composition is administered intramuscularly or subcutaneously as a general protection against 25 cancer development in internal organs.

27. The pharmaceutical composition of any of Claims 19 to 26 wherein the active fragment of lunasin peptide is selected from the group consisting of:

protein having amino acids 1 to 42 (SEQ. ID.2),


protein having amino acids 1 to 41 (SEQ. ID.3),

protein having amino acids 1 to 40 (SEQ. ID.4),
protein having amino acids 1 to 39 (SEQ. ID.5),
protein having amino acids 1 to 38 (SEQ. ID.6).
protein having amino acids 22 to 43 (SEQ. ID.7),
protein having amino acids 22 to 42 (SEQ. ID.8),
5 protein having amino acids 22 to 41 (SEQ. ID.9),
protein having amino acids 22 to 40 (SEQ. ID.10),
protein having amino acids 22 to 39 (SEQ. ID.11), and
protein having amino acids 22 to 38 (SEQ. ID.12),

28. The pharmaceutical composition of any of Claims 19 to 26 wherein the
10 therapeutic dose is about 250 microgram per milliliter or per gram of solid dose to about 2.5
milligram per milliliter or per gram of solid dose.

Expression Pattern of Lunasin

Stem
Root
Pod
Mature seed
Cotyledon



18S
rRNA
0.9 nt

18S
rRNA
0.9 nt

FIG. 1B


Expression Pattern of Lunasin

A Weeks after Flowering
Leaf 1 2 3 4 5 6 7



FIG. 1A


Immunolocalization of Lunasin



Lunasin immunostain
(arrows=Lunasin co-localizing
with endoreduplicated DNA)

FIG. 1E

Immunolocalization of Lunasin



Lunasin immunostain
(arrows=Lunasin co-localizing
with endoreduplicated DNA)

FIG. 1D


Immunolocalization of Lunasin



Histochemical stain
(arrows=endoreduplicated DNA)

FIG. 1C

2 / 13

**FIG. 2**




FIG._3C

FIG._3B

FIG._3A




FIG._3F




FIG._3E





FIG._3D

4 / 13



5 / 13



**FIG. 5****FIG. 6**

FIG. 7A

E1A-ΔCR1





FIG. 7B

E1A-ΔCR1 + Lunasin



FIG. 7C

E1A-wt



E1A-wt + Lunasin


FIG. 7D


Phase Contrast

FIG. 7E

Annexin V-FITC


FIG. 7F





Lunasin	S K W Q H Q Q Q D S C R K Q L Q G V N L T P C	E K H I M E K I	Q G R G D D D D D D D D D D
Lunasin-GRG	S K W Q H Q Q Q D S C R K Q L Q G V N L T P C	E K H I M E K I	Q D D D D D D D D D
trLunasin		E K H I M E K I	Q G R G D D D D D D D D D D
trLunasin-del		E K H I M E K I	Q G
NLS-trLunasin	L E E K Q K K K M E K E	Q G R G D D D D D D D D D D	

9 / 13

**FIG. 10**




FIG. 12A

Lunasin	S K W Q H Q Q D S C R K Q L Q G V N L T P C	E K H I M E K I	Q G R G D D D D D D D D D D
Lunasin-GRG	S K W Q H Q Q D S C R K Q L Q G V N L T P C	E K H I M E K I	Q D D D D D D D D D
trLunasin		E K H I M E K I	Q G R G D D D D D D D D D D
trLunasin-del		E K H I M E K I	Q G
NLS-trLunasin	L E E K Q K K K M E K E	Q G R G D D D D D D D D D D	D D D D D D D D D D

FIG. 12B

12 / 13





FIG. 13A

13 / 13

**FIG. 13B**

1/10

SEQUENCE LISTING

<110> FILGEN BIOSCIENCES, INC.

<120> THERAPEUTIC PEPTIDES HAVING A MOTIF THAT BINDS
SPECIFICALLY TO NON-ACETYLATED H3 AND H4 HISTONES FOR
CANCER THERAPY

<130> 3729.02-1PCT (HMP)

<140> PCT/US01/09453

<141> 2001-03-23

<150> 09/534,705

<151> 2000-03-24

<160> 28

<170> PatentIn Ver. 2.1

<210> 1

<211> 43

<212> PRT

<213> Glycine max

<400> 1

Ser Lys Trp Gln His Gln Gln Asp Ser Cys Arg Lys Gln Leu Gln Gly
1 5 10 15Val Asn Leu Thr Pro Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly
20 25 30Arg Gly Asp Asp Asp Asp Asp Asp Asp Asp
35 40

<210> 2

<211> 42

<212> PRT

<213> Glycine max

<400> 2

Ser Lys Trp Gln His Gln Gln Asp Ser Cys Arg Lys Gln Leu Gln Gly
1 5 10 15Val Asn Leu Thr Pro Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly
20 25 30Arg Gly Asp Asp Asp Asp Asp Asp Asp
35 40

<210> 3

<211> 41

<212> PRT

<213> Glycine max

2/10

<400> 3

Ser Lys Trp Gln His Gln Gln Asp Ser Cys Arg Lys Gln Leu Gln Gly
 1 5 10 15

Val Asn Leu Thr Pro Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly
 20 25 30

Arg Gly Asp Asp Asp Asp Asp Asp
 35 40

<210> 4

<211> 40

<212> PRT

<213> Glycine max

<400> 4

Ser Lys Trp Gln His Gln Gln Asp Ser Cys Arg Lys Gln Leu Gln Gly
 1 5 10 15

Val Asn Leu Thr Pro Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly
 20 25 30

Arg Gly Asp Asp Asp Asp Asp Asp
 35 40

<210> 5

<211> 39

<212> PRT

<213> Glycine max

<400> 5

Ser Lys Trp Gln His Gln Gln Asp Ser Cys Arg Lys Gln Leu Gln Gly
 1 5 10 15

Val Asn Leu Thr Pro Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly
 20 25 30

Arg Gly Asp Asp Asp Asp Asp
 35

<210> 6

<211> 38

<212> PRT

<213> Glycine max

<400> 6

Ser Lys Trp Gln His Gln Gln Asp Ser Cys Arg Lys Gln Leu Gln Gly
 1 5 10 15

Val Asn Leu Thr Pro Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly
 20 25 30

Arg Gly Asp Asp Asp Asp
 35

3/10

<210> 7
<211> 22
<212> PRT
<213> Glycine max

<400> 7
Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly Arg Gly Asp Asp Asp
1 5 10 15
Asp Asp Asp Asp Asp
20

<210> 8
<211> 21
<212> PRT
<213> Glycine max

<400> 8
Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly Arg Gly Asp Asp Asp
1 5 10 15
Asp Asp Asp Asp Asp
20

<210> 9
<211> 20
<212> PRT
<213> Glycine max

<400> 9
Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly Arg Gly Asp Asp Asp
1 5 10 15
Asp Asp Asp Asp
20

<210> 10
<211> 19
<212> PRT
<213> Glycine max

<400> 10
Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly Arg Gly Asp Asp Asp
1 5 10 15
Asp Asp Asp

<210> 11
<211> 18
<212> PRT
<213> Glycine max

4/10

<400> 11

Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly Arg Gly Asp Asp Asp
 1 5 10 15

Asp Asp

<210> 12

<211> 17

<212> PRT

<213> Glycine max

<400> 12

Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly Arg Gly Asp Asp Asp
 1 5 10 15

Asp

<210> 13

<211> 129

<212> DNA

<213> Glycine max

<400> 13

tccaaatggc agcaccagca agatagctgc cgcaagcagc tccaggggggt gaacctcacg 60
 ccctgcgaga agcacatcat ggagaagatc caaggccgcg gcgatgacga tgatgatgat 120
 gacgacgac 129

<210> 14

<211> 9408

<212> DNA

<213> Glycine max

<400> 14

agatctaaca tccaaagacg aaagggttgaa tgaaaccttt ttgccatccg acatccacag 60
 gtccattctc acacataagt gccaaaacgca acaggagggg atacactagc agcagaccgt 120
 tgcaaacgca ggacctccac tcctcttctc ctcaacacccc actttttgcca tcgaaaaacc 180
 agcccagtta ttgggcttga ttggagctcg ctcatccaa ttccttctat taggtacta 240
 acaccatgac tttattagcc tgtctatcct ggccccctg gcgaggttca tgtttgttta 300
 tttccgaatg caacaagctc cgcattacac ccgaacatca ctccagatga gggctttctg 360
 agtgtggggg caaatagttt catgttcccc aaatggccca aaactgacag tttaaacgct 420
 gtcttggaaac ctaatatgac aaaagcgtga tctcatccaa gatgaactaa gtttggttcg 480
 ttgaaatgct aacggccagt tgggtcaaaaa gaaacttcca aaagtcgcca taccgtttgt 540
 cttgttttgt attgattgac gaatgctcaa aaataatctc attaatgctt agcgcagtct 600
 ctctatcgct tctgaacccc ggtgcacctg tgccgaaaacg caaatgggga aacaccgct 660
 ttttggatga ttatgcattg tctccacatt gtatgcttcc aagattctgg tgggaatact 720
 gctgatagcc taacgttcat gatcaaaaatt taactgttct aaccctact tgacagcaat 780
 atataaacag aaggaagctg ccctgtctta aacctttttt tttatcatca ttattagctt 840
 actttcataa ttgcgactgg ttccaattga caagcttttg attttaacga cttttaacga 900
 caacttgaga agatcaaaaa acaactaatt attcgaagga tccaaacgat gagatttcct 960
 tcaattttta ctgcagtttt attcgcagca tcctccgcat tagctgctcc agtcaacact 1020
 acaacagaag atgaaacggc acaaattccg gctgaagctg tcategggta ctacagattta 1080
 gaaggggatt tcgatgttgc tgttttgcca ttttccaaca gcacaaataa cgggttattg 1140
 tttataaata ctactattgc cagcattgct gctaaagaag aaggggtatc tctcgagaaa 1200

agagaggctg	aagcttacgt	atccaaatgg	cagcaccagc	aagatagctg	ccgcaagcag	1260
ctccaggggg	tgaacctcac	gccctgcgag	aagcacatca	tggaagaagat	ccaaggccgc	1320
ggcgaatgacg	atgatgatga	tgaacgacgac	taagaattcc	ctagggcggc	cgcaattaa	1380
ttcgcccttag	acatgactgt	tcctcagttc	aagttgggca	cttacgagaa	gaccggtctt	1440
gctagattct	aatcaagagg	atgtcagaat	gccatttgcc	tgagagatgc	aggcttcatt	1500
tttgatactt	ttttatttgt	aacctatata	gtataggatt	ttttttgtca	ttttgtttct	1560
tctcgtacga	gcttgctcct	gatcagccta	tctcgcagct	gatgaatata	ttgtggtagg	1620
ggtttgggaa	aatcattcga	gtttgatgtt	tttcttggtg	ttccccactc	ctcttcagag	1680
tacagaagat	taagtgaaga	gttcgtttgt	gcaagcttat	cgataagctt	taatgcggta	1740
gtttatcaca	gttaaattgc	taacgcagtc	aggcaccgtg	tatgaaatct	aacaatgcgc	1800
tcacgtcat	cctcggcacc	gtcaccctgg	atgctgtagg	cataggcttg	gttatgccgg	1860
tactgccggg	cctcttgccg	gatatacgcc	attccgacag	catcgccagt	cactatggcg	1920
tgctgctagc	gctatatgcg	ttgatgcaat	ttctatggcg	accggttctc	ggagcaactgt	1980
ccgaccgctt	tgcccgccgc	ccagtcctgc	tcgcttcgct	acttgagacc	actatcgact	2040
acgcgatcat	ggcgaccaca	cccgctcctgt	ggatctatcg	aatctaaatg	taagttaaaa	2100
tctctaaata	attaaataag	tcccagtttc	tccatacgaa	ccttaacagc	attgcggtga	2160
gcacttagac	cttcaacagc	agccagatcc	atcactgctt	ggccaatatg	tttcagttcc	2220
tcaggagtta	cgtcttgtga	agtgatgaac	ttctggaagg	ttgcagtgtt	aactccgctg	2280
tattgacggg	catatccgta	cgttggcaaa	gtgtgggttg	taccggagga	gtaatctcca	2340
caactctctg	gagagtaggc	accaacaaac	acagatccag	cgtgttgtag	ttgatcaaca	2400
taagaagaag	cattctcgat	ttgcaggatc	aagtgttcag	gagcgtactg	attggacatt	2460
tccaaagcct	gctcgtaggt	tgcaaccgat	agggttgtag	agtgtgcaat	acacttgctg	2520
acaatttcaa	cccttggaac	ctgcacagct	tggttgtaga	cagcatcttc	aattctggca	2580
agctccttgt	ctgtcatatc	gacagccaac	agaatcacct	gggaatcaat	accatgttca	2640
gcttgagaca	gaaggctctga	ggcaacgaaa	tctggatcag	cgtatttatc	agcaataact	2700
agaacttcag	aaggccagc	aggcatgtca	atactacaca	gggctgatgt	gtcattttga	2760
accatcatct	tggcagcagt	aacgaactgg	ttcctggac	caaataattt	gtcacactta	2820
ggaacagttt	ctgttccgta	agccatagca	gctactgcct	gggcgcctcc	tgctagcacg	2880
atacacttag	caccaacctt	gtgggcaacg	tagatgactt	ctggggtaag	ggtaccatcc	2940
ttcttaggtg	gagatgcaaa	aacaatttct	ttgcaaccag	caactttggc	aggaacaccc	3000
agcatcaggg	aagtggaaag	cagaatttgcg	gttccaccag	gaatatagag	gccaactttc	3060
tcaataggtc	ttgcaaaaacg	agagcagact	acaccagggc	aagtctcaac	ttgcaacgtc	3120
tccgttagtt	gagcttcatg	gaatttccctg	acgttatcta	tagagagatc	aatggctctc	3180
ttaacgttat	ctggcaattg	cataagttcc	tctgggaaag	gagcttctaa	cacagggtgtc	3240
ttcaaagcga	ctccatcaaa	cttggcagtt	agttctaaaa	gggctttgtc	accattttga	3300
cgaaacattgt	cgacaatttg	tttgactaat	tccataatct	gttccgtttt	ctggatagga	3360
cgacgaaggg	catcttcaat	ttcttgtgag	gaggccttag	aaacgtcaat	tttgcaaat	3420
tcatacgcag	cttcgaagg	gacttcttta	ggtttggatt	cttctttagg	ttgttccttg	3480
gtgtatcctg	gcttggcatc	tcctttcctt	ctagtgcact	ttagggactt	catatccagg	3540
tttctctcca	cctcgtccaa	cgtcacaccg	tacttggcac	atctaactaa	tgcaaaataa	3600
aataagtcag	cacattccca	ggctatatct	tccttggatt	tagcttctgc	aagttcatca	3660
gcttcctccc	taatttttagc	gttcaacaaa	acttcgtcgt	caaataaccg	tttggataaa	3720
gaaccttctg	gagcattgct	cttacgatcc	cacaagggtg	cttccatggc	tctaagaccc	3780
tttgattggc	caaaacagga	agtgcgttcc	aagtgcagca	aaccaacacc	tgtttgttca	3840
accacaaatt	tcaagcagtc	tccatcacaa	tccaattcga	taccagcaa	cttttgagtt	3900
gtccagatg	tagcaccttt	ataccacaaa	ccgtgacgac	gagattggta	gactccagtt	3960
tgtgtcctta	tagcctccgg	aatagacttt	ttggacgagt	acaccaggcc	caacgagtaa	4020
ttagaagagt	cagccaccaa	agtagtgaat	agaccatcgg	ggcggtcagt	agtcaaagac	4080
gccaacaaaa	tttactgac	agggaacttt	ttgacatctt	cagaaagttc	gtattcagta	4140
gtcaattgcc	gagcatcaat	aatggggatt	ataccagaag	caacagtggg	agtcacatct	4200
accaactttg	cggctctcaga	aaaagcataa	acagttctac	taccgccatt	agtgaacttt	4260
ttcaaactcg	ccagtggaga	agaaaaaggc	acagcgatac	tagcattagc	gggcaaggat	4320
gcaactttat	caaccagggt	cctatagata	accctgacgc	ctgggatcat	cctttggaca	4380
actctttctg	ccaaaactag	gtccaaaatc	acttcattga	taccattatt	gtacaacttg	4440
agcaagttgt	cgatcagctc	ctcaaattgg	tcctctgtaa	cggatgactc	aacttgcaca	4500
ttaaacttgaa	gctcagtcga	ttgagtgaac	ttgatcaggt	tgtgcagctg	gtcagcagca	4560
tagggaaaca	cggcttttcc	taccaaactc	aagggaattat	caaactctgc	aacacttgcg	4620
tatgcaggta	gcaaggga	tgtcatactt	gaagtcggac	agtgagtgtg	gtcttgagaa	4680

attctgaagc cgtatttttta ttatcagtga gtcagtcac aggagatcct ctacgccgga 4740
cgcatcgtgg ccgacctgca gggggggggg gggcgctgag gtctgcctcg tgaagaaggt 4800
gttgcctgact cataccaggc ctgaatcgcc ccatcatcca gccagaaagt gagggagcca 4860
cgggtgatga gagctttgtt gtaggtggac cagttggtga ttttgaactt ttgctttgcc 4920
acggaacggt ctgctgtgtc ggggaagatgc gtgatctgat ccttcaactc agcaaaagt 4980
cgattttattc aacaaagccg ccgtcccgtc aagtcagcgt aatgctctgc cagtggtaca 5040
accaattaac caattctgat tagaaaaact catcgagcat caaatgaaac tgcaatttat 5100
tcataatcagg attatcaata ccatattttt gaaaaagccg tttctgtaat gaaggagaaa 5160
actcaccgag gcagttccat aggatggcaa gatcctggta tcgggtctgcg attccgactc 5220
gtccaacatc aatacaacct attaatctcc cctcgtcaaa aataagggtta tcaagtgaga 5280
aatcaccatg agtgacgact gaatccgggtg agaattggcaa aagcttatgc atttctttcc 5340
agacttggtc aacaggccag ccattacgct cgtcatcaaa atcactcgca tcaaccaaac 5400
cgttattcat tcgtgattgc gcctgagcga gacgaaatac gcgatcgctg ttaaaaggtac 5460
aattacaaac aggaatcgaa tgcaaccggc gcagggaacac tgccagcgca tcaacaatat 5520
tttcacctga atcaggatat tcttctaata cctggaatgc tgttttcccg gggatcgag 5580
tggtgagtaa ccattgcatc tcaggagtag ggataaaatg cttgatggtc ggaagaggca 5640
taaattccgt cagccagttt agtctgacca tctcatctgt aacatcattg gcaacgctac 5700
ctttgccatg tttcagaaac aactctggcg catcgggctt cccatacaat cgatagattg 5760
tcgcacctga ttgcccgaca ttatcgcgag cccattttata cccatataaa tcagcatcca 5820
tggttggaatt taatcgcggc ctcgagcaag acgtttcccg ttgaatatgg ctcataaacac 5880
cccttgatt actgtttatg taagcagaca gttttattgt tcatgatgat atatttttat 5940
cttggtgcaat gtaacatcag agattttgag acacaacgtg gctttcccc cccccctgc 6000
aggctggcat caccggcgcc acagggtcgg ttgctggcgc ctatatcgcc gacatcaccg 6060
atgggggaaga tcgggctcgc cacttcgggc tcatgagcgc ttgtttcggc gtgggtatgg 6120
tggcagggccc cgtggccggg ggactgttgg gcgccatctc cttgcatgca ccattccttg 6180
cggcgcgcggt gctcaacggc ctcaacctac tactgggctg cttcctaatt caggagtcgc 6240
ataagggaga gcgtcgagta tctatgattg gaagtatggg aatgggtgata cccgcattct 6300
tcagtgtctt gaggtctcct atcagattat gcccaactaa agcaaccgga ggaggagatt 6360
tcattgtaaa tttctctgac ttttggatc cagtagactc gaactgtgag actatctcg 6420
ttatgacagc agaaatgtcc ttcttgga cagtaaatga agtccacca ataaagaaat 6480
ccttggtatc aggaacaaac ttctgtttc gaactttttc ggtgccttga actataaaat 6540
gtagagtggg tatgtcgggt aggaatggag cgggcaaatg cttaccttct ggaccttcaa 6600
gaggtatgta ggggtttag atactgatgc caacttcagt gacaacggtg ctatttcgtt 6660
caaaccattc cgaatccaga gaaatcaaag ttgtttgtct actattgatc caagccagt 6720
cggctctgaa actgacaata gtgtgctcgt gttttgaggt catctttgta tgaataaact 6780
tagtctttga tctaaataat cttgacgagc caaggcgata aatacccaaa tctaaaactc 6840
ttttaaaacg ttaaaaggac aagtatgtct gctgtatta aaccccaaat cagctcgtag 6900
tctgatcttc atcaacttga ggggcactat cttgttttag agaaatttgc ggagatgcga 6960
tatcgagaaa aaggtagcgt gatttttaaac gtgaaattta tctcaagatc tctgcctcgc 7020
geggttcggt gatgacggtg aaaacctctg acacatgcag ctcccgagga cggtcacagc 7080
ttgtctgtaa ggggatgccc ggagcagaca agcccgtcag ggcgctcag cgggtgttgg 7140
cgggtgtcgg ggcgcagcca tgaccagtc acgtagcgat agcggagtgt atactggctt 7200
aactatgccc catcagagca gattgtactg agagtgcacc atatgcggtg tgaaataccg 7260
cacagatgcg taaggagaaa ataccgcatc aggcgctctt cctactcaaa ggcggttaata 7380
tcgctgctcgt cggctgttcg gctgcgcca ggataacgca ggaaagaaca tgtgagcaaa aggcagcaa 7440
aaggccagga accgtaaaaa ggccgcgttg ctggcgtttt tccataggct ccgccccct 7500
gacgagcatc acaaaaatcg acgctcaagt cagaggtggc gaaaccgac aggactataa 7560
agataccagg cgtttccccc tggaagctcc ctcgtgcgt ctcctgttcc gacctgcg 7620
cttaccggat acctgtccgc ctttctccct tcgggaagcg tggcgcttcc tcaatgctca 7680
cgctgtagggt atctcagttc ggtgtaggtc gttcgtcca agctgggctg tgtgcacgaa 7740
cccccggttc agcccgaccg ctgcccctta tccggttaact atcgtcttga gtccaaccg 7800
gtaagacagc acttatcgcc actgpcagca gccactggta acaggattag cagagcgagg 7860
tatgtaggcg gtgctacaga gttcttgaag tgggtggccta actacggcta cactagaagg 7920
acagtatttg gtatctgcgc tctgtgaag ccagttacct tcggaaaaag agttggtagc 7980
tcttgatccg gcaaaaaaac caccgctgggt agcgggtgggt tttttgtttg caagcagcag 8040
attacgcgca gaaaaaaagg atctcaagaa gatcctttga tcttttctac ggggtctgac 8100
gctcagtggg acgaaaactc acgttaaggg attttggta tgagattatc aaaaaggatc 8160

7/10

```

ttcacctaga tcctttttaa ttaaaaatga agtttttaaat caatctaaag tatatatgag 8220
taaacttggg ctgacagtta ccaatgctta atcagtgagg cacctatctc agcgatctgt 8280
ctatttcgtt catccatagt tgcctgactc cccgtcgtgt agataactac gatacgggag 8340
ggcttaccat ctggcccccag tgctgcaatg ataccgcgag acccacgctc accggctcca 8400
gatttatcag caataaacca gccagccgga agggccgagc gcagaagtgg tcctgcaact 8460
ttatccgcct ccatccagtc tattaattgt tgccgggaag ctagagtaag tagttcgcca 8520
gttaaatagt tgcgcaacgt tggtgccatt gctgcaggca tcgtggtgtc acgctcgtcg 8580
tttggtatgg cttcattcag ctccggttcc caacgatcaa ggcgagttac atgatcccc 8640
atgttggtga aaaaagcggg tagctccttc ggtcctccga tcgttggtcag aagtaagttg 8700
gccgcagtgt tatcactcat gggtatggca gcactgcata attctcttac tgtcatgcc 8760
tccgtaagat gcttttctgt gactggtgag tactcaacca agtcattctg agaatagtgt 8820
atgcccgcac cgagttgctc ttgcccggcg tcaacacggg ataataccgc gccacatagc 8880
agaactttaa aagtgtcat cattggaaaa cgttcttcgg ggcgaaaact ctcaaggatc 8940
ttaccgctgt tgagatccag ttcgatgtaa cccactcgtg caccactg atcttcagca 9000
tccttttact tcaccagcgt ttctgggtga gcaaaaacag gaaggcaaaa tgccgcaaaa 9060
aagggaataa gggcgacacg gaaatgttga atactcatac tcttcctttt tcaatattat 9120
tgaagcattt atcaggggta ttgtctcatg agcggataca tatttgaatg tathtagaaa 9180
aataaaca aa taggggttcc gcgcacattt ccccgaaaag tgccaccta cgtctaagaa 9240
accattatta tcatgacatt aacctataaa aataggcgta tcacgaggcc ctttcgtctt 9300
caagaattaa ttctcatgtt tgacagctta tcatcgataa gctgactcat gttgggtattg 9360
tgaaatagac gcagatcggg aacactgaaa aataacagtt attattcg 9408

```

<210> 15

<211> 40

<212> PRT

<213> Glycine max

<220>

<223> Lunasin-GRG

<400> 15

```

Ser Lys Trp Gln His Gln Gln Asp Ser Cys Arg Lys Gln Leu Gln Gly
1          5          10          15

```

```

Val Asn Leu Thr Pro Cys Glu Lys His Ile Met Glu Lys Ile Gln Asp
20          25          30

```

```

Asp Asp Asp Asp Asp Asp Asp Asp
35          40

```

<210> 16

<211> 21

<212> PRT

<213> Glycine max

<220>

<223> trLunasin

<400> 16

```

Glu Lys His Ile Met Glu Lys Ile Gln Gly Arg Gly Asp Asp Asp Asp
1          5          10          15

```

```

Asp Asp Asp Asp Asp
20

```

8/10

<210> 17
<211> 10
<212> PRT
<213> Glycine max

<220>
<223> trLunasin-del

<400> 17
Glu Lys His Ile Met Glu Lys Ile Gln Gly
1 5 10

<210> 18
<211> 25
<212> PRT
<213> Glycine max

<220>
<223> NLS-trLunasin

<400> 18
Leu Glu Glu Lys Gln Lys Lys Lys Met Glu Lys Glu Gln Gly Arg Gly
1 5 10 15

Asp Asp Asp Asp Asp Asp Asp Asp
20 25

<210> 19
<211> 11
<212> PRT
<213> Glycine max

<220>
<223> Lunasin

<400> 19
Cys Glu Lys His Ile Met Glu Lys Ile Gln Gly
1 5 10

<210> 20
<211> 12
<212> PRT
<213> Glycine max

<220>
<223> HuHP1 (p25)

<400> 20
Glu Glu Glu Glu Tyr Val Val Glu Lys Val Leu Asp
1 5 10

<210> 21
<211> 12
<212> PRT

9/10

<213> Glycine max

<220>

<223> DmPc

<400> 21

Val Asp Leu Val Tyr Ala Ala Glu Lys Ile Ile Gln
1 5 10

<210> 22

<211> 12

<212> PRT

<213> Glycine max

<220>

<223> Hu HP1B

<400> 22

Phe Glu Arg Gly Leu Glu Pro Glu Lys Ile Ile Gly
1 5 10

<210> 23

<211> 12

<212> PRT

<213> Glycine max

<220>

<223> SpSw16A

<400> 23

Glu Glu Asp Glu Tyr Val Val Glu Lys Val Leu Lys
1 5 10

<210> 24

<211> 12

<212> PRT

<213> Glycine max

<220>

<223> PcHET2A

<400> 24

Val Glu Glu Glu Phe Ile Val Glu Lys Ile Leu Asp
1 5 10

<210> 25

<211> 12

<212> PRT

<213> Glycine max

<220>

<223> DvHP1A

10/10

<400> 25

Glu Glu Glu Glu Tyr Ala Val Glu Lys Ile Leu Asp
1 5 10

<210> 26

<211> 12

<212> PRT

<213> Glycine max

<220>

<223> MoMOD1A

<400> 26

Glu Glu Glu Glu Tyr Val Val Glu Lys Val Leu Asp
1 5 10

<210> 27

<211> 12

<212> PRT

<213> Glycine max

<220>

<223> SmPAT26

<400> 27

Gly Glu Asp Glu Phe Gln Val Glu Lys Ile Leu Lys
1 5 10

<210> 28

<211> 12

<212> PRT

<213> Glycine max

<220>

<223> DmHP1A

<400> 28

Glu Glu Glu Glu Tyr Ala Val Glu Lys Ile Ile Asp
1 5 10

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number
WO 01/72784 A3

(51) International Patent Classification⁷: **C07K 14/415**,
C12N 15/29, A61K 38/16, A61P 35/00

[PH/US]; 2800 Dover Avenue #G-5, Fairfield, CA 94533 (US).

(21) International Application Number: PCT/US01/09453

(74) Agents: **PETERS, Howard, M.** et al.; Peters, Verny, Jones & Biksa, LLP, Suite 6, 385 Sherman Avenue, Palo Alto, CA 94306 (US).

(22) International Filing Date: 23 March 2001 (23.03.2001)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:
09/534,705 24 March 2000 (24.03.2000) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/534,705 (CIP)
Filed on 24 March 2000 (24.03.2000)

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **FILGEN BIOSCIENCES, INC.** [US/US]; 2333 Courage Drive, Suite C, Fairfield, CA 94533 (US).

(72) Inventor; and


Published:

(75) Inventor/Applicant (*for US only*): **GALVEZ, Alfredo, F.**

— with international search report

[Continued on next page]

(54) Title: **PEPTIDES BINDING TO NON-ACETYLATED H3 AND H4 HISTONES FOR CANCER THERAPY**



(57) Abstract: The present invention describes a composition of matter comprising a conserved structural motif that allows the targeting and binding of a chromatin binding protein to non-acetylated histone H3 and H4 and prevents their acetylation. This invention is responsible for the anti-carcinogenic property of a chromatin binding peptide isolated from soybean seed. This structural motif is found in a highly conserved manner in other chromatin-binding proteins from different species. Modifications to this structural motif such as fusions to other proteins with functional motifs and amino acid substitutions have potential therapeutic applications and can be developed as an *in vivo* gene silencing technology for biological and medical research. In particular, active fragments of the lunasin peptide and active analogs of the lunasin peptide are useful in this invention. Pharmaceutical compositions useful in retarding or stopping or reducing various types of cancers are described.

WO 01/72784 A3



(88) Date of publication of the international search report:
6 June 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/09453

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/415 C12N15/29 A61K38/16 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, PAJ, SEQUENCE SEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 15642 A (UNIV CALIFORNIA) 1 April 1999 (1999-04-01) cited in the application page 1, line 27 -page 3, line 10 examples 4,5	1-28
A	--- GALVEZ A F: "A soybean cDNA encoding a chromatin-binding peptide inhibits mitosis of mammalian cells" NATURE BIOTECHNOLOGY, NATURE PUBLISHING, US, vol. 17, no. 5, May 1999 (1999-05), pages 495-500, XP002170660 ISSN: 1087-0156 the whole document --- -/--	1-28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

*Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

14 March 2002

Date of mailing of the international search report

21/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

De Kok, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/09453

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 00 66625 A (UNIV CALIFORNIA) 9 November 2000 (2000-11-09) the whole document	1-28
P,X	LAM Y ET AL: "IN VITRO AND IN VIVO CHEMOPREVENTIVE PROPERTIES OF A SOYBEAN PEPTIDE (LUNASIN)" FASEB JOURNAL (FEDERATION OF AMERICAN SOCIETIES FOR EXPERIMENTAL BIOLOGY), BETHESDA, US, vol. 15, no. 4, 7 March 2001 (2001-03-07), page A281 XP001025557 ISSN: 0892-6638 abstract	1-28
E	WO 01 34808 A (FILGEN BIOSCIENCES INC) 17 May 2001 (2001-05-17) the whole document	1-28

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1, 6, 8, 13, 15, 17, all partially

Present claims 1, 6, 8, 13, 15 and 17 relate to lunasin fragments or analogs defined by reference to a desirable characteristic or property, namely by being "active".

The claims cover all fragments/analogues having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such fragments/analogues. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the fragments/analogues by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the native lunasin peptide and to the fragments/analogues identified in claim 7 resp. 16 and 27.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/09453

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9915642	A	01-04-1999	US 6107287 A	22-08-2000
			AU 731631 B2	05-04-2001
			AU 9582198 A	12-04-1999
			CA 2303061 A1	01-04-1999
			EP 1017798 A1	12-07-2000
			JP 2001517434 T	09-10-2001
			WO 9915642 A1	01-04-1999
WO 0066625	A	09-11-2000	AU 4809600 A	17-11-2000
			EP 1173476 A1	23-01-2002
			WO 0066625 A1	09-11-2000
WO 0134808	A	17-05-2001	AU 1918301 A	06-06-2001
			WO 0134808 A2	17-05-2001

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.